

# **PIPE LAKE 2017**

## **Lake Stewardship Monitoring Report**

King County Water & Land Resources Division  
Science & Technical Support Section  
[www.kingcounty.gov/EnvironmentalScience](http://www.kingcounty.gov/EnvironmentalScience)

---

### **Summary & Recommendations**

*Thank you to Tom Sullivan, the volunteer monitor for Pipe Lake*

**The key takeaways from the 2017 monitoring season are:**

- Pipe Lake continued to have clear water, low nutrient concentrations, and low algal growth.
- Pipe Lake has one of the deepest Secchi depths (clearest water) of any lake in the Lake Stewardship program. Secchi depths have been getting even deeper over time.
- No algal blooms were reported in 2017.

**The Lake Stewardship Program recommends:**

- Monitoring is a key part of good lake stewardship, building a valuable long-term dataset to guide lake management and detect any future problems. Continue to monitor Pipe Lake through the Lake Stewardship Program.
- 

### ***In this report:***

- **What We Measure & Why**
- **Water Quality Results & Trends**
  - 2017 monitoring results
  - Long-term annual averages
  - Trends over time
- **Trophic State**
  - Trophic state indices
  - Comparison map
- **Supplemental Data**
  - Summary statistics
  - Water column profile
  - Total alkalinity
  - Water color

## **What We Measure & Why**

- **Secchi depth** is a measure of water clarity or transparency. Secchi depth is shallower when there are more suspended particles in the lake, such as sediment or algae. Secchi depth is also affected by water color, often from tannins or other naturally occurring organic molecules.
- **Water temperature** can affect the growth rates of plants and algae. In addition, cooler or warmer water temperatures favor different species of fish and other aquatic organisms.
- **Chlorophyll-a** is a measure of the amount of algae in a lake. Chlorophyll-a is a pigment necessary for algae to photosynthesize and store energy.
- **Phosphorus** and **nitrogen** are naturally occurring nutrients necessary for growth and reproduction in both plants and animals. Increases in nutrients (especially phosphorus) can lead to more frequent and dense algal blooms.
- The **ratio of total nitrogen to total phosphorus (N:P)** indicates whether nutrient conditions favor the growth of cyanobacteria (blue-green algae). When N:P ratios are near or below 25, cyanobacteria can dominate the algal community. This is important because cyanobacteria can potentially produce toxins.


## **Water Quality Results & Trends**

The following graphs show the water-quality parameters that are sampled from May through October, at 1 m depth (additional depths and parameters are measured on profile days; see *Supplemental Data*). The left column of graphs shows results for each sampling date in 2017, and the right column shows average values for each year (May-October averages).


Data for Pipe Lake are the blue circles (with white centers) connected by the blue line. Any gaps in the blue line indicate missed samples. To provide some context for these values, the grey points in the background are results for all other lakes in the Lake Stewardship program.

Any long-term trends in Pipe Lake are drawn with a dashed red line and described further after the graphs. Statistical trend analyses used a seasonal (monthly) Kendall test ( $p < 0.05$ ).

## 2017 Monitoring Results



## Long-Term Annual Averages



The table below gives more details about the long-term trends. Results are presented as an average amount and percent of change per decade (the increase or decrease over ten years). Percent change is calculated as the percent of the estimated value in 1994, when monitoring started.

| Parameter         | Change per Decade | (%)     |
|-------------------|-------------------|---------|
| Secchi depth      | 0.91 m            | (23%)   |
| Water temperature | 0.9 °C            | (5.1%)  |
| Total nitrogen    | -26 µg/L          | (-6.1%) |
| N:P ratio         | -5.8              | (-12%)  |


The trend of deeper Secchi depths (clearer water) in Pipe Lake is especially noticeable. On average, Secchi depths in recent years have been about 1.5 m deeper than they were in the mid-1990s when monitoring began.

## **Trophic State**

The Trophic State Index (TSI) is a common index of a lake's overall biological productivity. TSI values are calculated from Secchi depth, chlorophyll-a concentrations, and total phosphorus concentrations. These three TSI estimates are all scaled between 0 and 100.

*Oligotrophic* lakes (TSI <40) are very clear, with low nutrient concentrations and low algal growth. *Eutrophic* lakes (TSI >50) have cloudy water, with high nutrient concentrations and high algal growth. Eutrophic lakes are more likely to have frequent algal blooms. *Mesotrophic* lakes (TSI 40-50) are in the middle, with fairly clear water, and moderate nutrient concentrations and algal growth. Lakes in lowland King County have a range of different natural trophic states, and human activities may also alter a lake's trophic state (usually by changing nutrient inputs).


## **Trophic state indices**



In recent years, the three TSI values have been in the oligotrophic range, or near the oligotrophic-mesotrophic boundary.

## Comparison map

For a comparison with other lakes, this map shows the trophic state for each lake in the King County Lake Stewardship program in 2017. The color of each circle indicates the lake's average chlorophyll-a TSI value for the year.



## Supplemental Data

### Summary statistics

This table summarizes data from May-October 2017 (1 m depth only), giving the minimum, mean (average), and maximum values for each parameter. To reduce biases from missing data or changes in sampling frequency, monthly means were calculated and then averaged to give an overall mean.

| Parameter               | Minimum | Mean  | Maximum |
|-------------------------|---------|-------|---------|
| Secchi depth (m)        | 5.3     | 6.3   | 7.0     |
| Water temperature (°C)  | 15.5    | 21.0  | 26.5    |
| Chlorophyll-a (µg/L)    | 0.8     | 1.8   | 3.3     |
| Total nitrogen (µg/L)   | 263.0   | 330.0 | 484.0   |
| Total phosphorus (µg/L) | 8.2     | 11.8  | 17.8    |
| N:P ratio               | 22.0    | 28.8  | 36.1    |

### Water column profile

In May and August, water was collected at the mid-lake sampling station from three depths in a water-column profile: 1 m, the middle depth of the water column, and 1 m from the lake bottom.

| Date      | Depth | Temp | Chlor | Pheo  | TN   | NH3    | NO2/3  | TP    | OPO4 |
|-----------|-------|------|-------|-------|------|--------|--------|-------|------|
| 5/22/2017 | 1     | 21.0 | 0.8   | (1.2) | 377  | 20.8   | 136.0  | 11.7  | 1.1  |
|           | 10    | 9.0  | 7.6   | 1.7   | 427  | –      | –      | 12.0  | –    |
|           | 19    | 9.0  | –     | –     | 717  | 40.3   | 389.0  | 92.5  | 16.9 |
| 8/22/2017 | 1     | 24.0 | 1.5   | (1.2) | 310  | 8.5    | 10.0   | 9.1   | 0.6  |
|           | 10    | 9.0  | 3.0   | 2.5   | 480  | –      | –      | 15.8  | –    |
|           | 19    | 7.0  | –     | –     | 1410 | 1130.0 | (10.0) | 418.0 | 93.3 |


\* Parameter abbreviations are: chlorophyll-a (Chlor), pheophytin (Pheo), total nitrogen (TN), ammonia (NH3), nitrate/nitrite (NO2/3), total phosphorus (TP), orthophosphate (OPO4). Depth is in m, temperature is in °C, and all other parameters are in µg/L.

Dashes indicate parameters that were not analyzed for a given sample. Values below the method detection limit (MDL) are enclosed in parentheses and have the value of the MDL substituted.

## Total alkalinity

A lake's ability to resist acidification, also called its buffering capacity, is measured as "total alkalinity." Lakes with total alkalinity less than 20 mg CaCO<sub>3</sub> are considered sensitive to acidification. We measured total alkalinity in May and August (on profile-sampling days) at 1 m depth. In 2017, the average total alkalinity of these two samples was 27.6 mg CaCO<sub>3</sub>.

The blue circles (with white centers) and blue line are annual average alkalinity values for Pipe Lake. Grey points in the background are results for all other lakes in the Lake Stewardship program. The dashed red line shows the long-term trend in alkalinity, with an average change of -5.48 mg CaCO<sub>3</sub> per decade.






## Water color

Water color affects a lake's water clarity (and Secchi depth). Water color is measured by shining a specific wavelength of ultraviolet light (254 nm) through a filtered water sample and measuring the percent that was absorbed. We measured UV254 absorbance in May and August (on profile-sampling days) at 1 m depth. In 2017, the average UV254 absorbance of these two samples was 0.08, on a scale where 0 is no absorbance (perfectly clear) and 1 is complete absorbance (perfectly opaque).

The blue circles (with white centers) and blue line are annual average UV absorbance values for Pipe Lake. Grey points in the background are results for all other lakes in the Lake Stewardship program.



### King County

Department of Natural Resources and Parks  
Water and Land Resources Division  
**Science and Technical Support Section**  
King Street Center, KSC-NR-0600  
201 South Jackson Street, Suite 600  
Seattle, WA 98104  
206-477-4800 TTY Relay: 711  
[www.kingcounty.gov/EnvironmentalScience](http://www.kingcounty.gov/EnvironmentalScience)